


PINEAPPLE

CADANGAN PENABURAN BAJA REALSTRONG UNTUK TANAMAN NANAS

RECOMMENDATION FOR REALSTRONG FERTILIZER APPLICATION FOR PINEAPPLE PLANTS



MASA PEMBAJAJAN / FERTILIZATION TIME	JENIS BAJA & KADAR / TYPE OF FERTILIZER & RATE	CATATAN/ REMARKS
SEBELUM MENANAM / BEFORE PLANTING	K-NEUTRALIZER 1KG/M ²	Disarankan untuk menabur K-Neutralizer di atas nutrien K dan elemen lain. It is recommended to spread K-Neutralizer on top of potassium and other nutrients.
SELEPAS 2 BULAN DITANAM / AFTER 2 MONTHS OF PLANTING	SOIL CONDITIONER 20-30G/LUBANG	
SELEPAS 4 BULAN DITANAM / AFTER 4 MONTHS OF PLANTING	OR MAX GROWTH 20-30G/LUBANG	
SELEPAS 6-7 BULAN DITANAM / AFTER 6-7 MONTHS OF PLANTING	MAX K 30G/LUBANG	-
SELEPAS 8 BULAN DITANAM / AFTER 8 MONTHS OF PLANTING	HORMON ETHREL	Jika pokok nanas subur, hormon Ethrel boleh digunakan. Tetapi, elakkan dari bercampur dengan baja. Oleh itu, hormon ini hanya boleh digunakan setelah 2 bulan dari hari menabur baja. If the pineapple plants are healthy, Ethrel hormone can be used. However, avoid mixing it with fertilizer. Therefore, this hormone should only be used after 2 months from the day of fertilization.
MUSIM BERBUAH (150 HARI) / FRUITING SEASON (150 DAYS)	MAX K 30G/LUBANG	-
2 KALI SEBULAN / 2 TIMES PER MONTH	TE + CaO, B, MgO 1L/HA (SETIAP KALI/EVERY TIME)	KAEDAH SEMBURAN / SPRAYING METHOD

Disediakan oleh Dr Noor Aisyah Azman (Agriculture Technical Executive, ACI).

